

OXFORD COUNTY KEY FINDINGS

- 2008 Population Estimate = 56,608
- 2008 Median Household Income 2008 = \$38,987
- 16% of residents are age 65+
- 30% of residents enrolled in Medicaid

Health Risks and Challenges

Health Assets and Opportunities

Risk Factors

- **Access to Care:**
 - High percentage of nonelderly adults (18-64) uninsured [OXF=18%, ME =16%]
 - Highest percentage of females with no usual source of medical care [OXF=12%, ME=8%]
 - High percentage reporting unmet medical needs in past year due to cost [OXF=9%, ME=7%]
 - High percentage with no dental visit in past 2 years [OXF=34%, ME=24%]
- **Health Status:**
 - High percentage report health as fair to poor [OXF=23%, ME=15%]
 - High percentage report 11 or more days lost in past month due to poor health [OXF=11%, ME=8%]
 - High percentage with 3+ chronic conditions [OXF=17%, ME=13%]
- **Overweight/Obesity:**
 - High percentage with sedentary lifestyle [OXF=24%, ME=21%]
 - High percentage obese [OXF=33%, ME=28%]
- **Smoking:** High percentage of current smokers: [OXF=25%, ME=22%], especially among males
- **Interpersonal violence:** High percentage experienced interpersonal violence in lifetime [OXF=15%, ME=12%]
- **Alcohol and Substance Use:**
 - High percentage of chronic heavy drinking [OXF=8%, ME=6%]
 - High percentage report use of street drugs in past 30 days [OXF=7%, ME=5%] and household overdose in past year [OXF=1.5%, ME=0.9%]
- **Youth (Grades 9-12):**
 - High percentage past month smokeless tobacco use [OXF=11%, ME=9.5%]
 - Low percentage report regular physical activity [OXF=34%, ME=39%]
 - High percentage overweight [OXF=17%, ME=14%] and obese [OXF=16%, ME=13%]

- **Smoking:** High percentage of smokers report having tried to quit in past year [OXF=61%, ME=54%] and having ever used Maine Tobacco Quitline [OXF=22%, ME=19%]
- **Prevention:** High percentage of elderly females report having received pneumococcal vaccination [OXF=87%, ME=74%]
- **Alcohol and Substance Use:** Low percentage with prescription drug misuse in past 30 days [OXF=0.9%, ME=1.8%]
- **Reproductive Health:** Low teen birth rate

Health Risks and Challenges

Health Assets and Opportunities

Disease Incidence & Prevalence

- | | |
|---|--|
| <ul style="list-style-type: none"> • Respiratory Disease: High prevalence of COPD [OXF=6.7%, ME=4.2%] and youth asthma [OXF=7%, ME=6%] • Cancer: <ul style="list-style-type: none"> ○ High percentage reporting ever diagnosed with cancer [OXF=9.3%, ME=7.5%] ○ High incidence of lung cancer among males ○ High incidence of bladder and colorectal cancers • Diabetes: <ul style="list-style-type: none"> ○ High prevalence of diabetes [OXF=12%, ME=10%] ○ Low receipt of hemoglobin A1c testing and foot examinations among diabetics • Infectious Disease: High incidence of gonorrhea • Mental Health: <ul style="list-style-type: none"> ○ High percentage reporting 11 or more days mental health not good in past month [OXF=14%, ME=11%] ○ High percentage with unmet mental health care needs [OXF=6%, ME=5%] ○ Highest percentage of any county considered at risk for depression based on MHI-5 score [OXF=12%, ME=7%] | <ul style="list-style-type: none"> • Cancer: Low incidence of female cervical cancer |
|---|--|

Hospital Utilization & Mortality Rates

- | | |
|---|--|
| <ul style="list-style-type: none"> • Hospital Admissions: <ul style="list-style-type: none"> ○ High hospitalization rates for respiratory disease including COPD, pneumonia, emphysema and bronchitis/asthma ○ High hospitalization rates for drug related psychoses, high risk pregnancy complications, hip procedures and head/brain injuries • Emergency Department (ED) Visits: <ul style="list-style-type: none"> ○ High overall ED use ○ High ED rates for bronchitis/asthma among youth (0-17) ○ High ED visit rates for COPD • Mortality: <ul style="list-style-type: none"> ○ High cerebrovascular (stroke) and COPD mortality rates ○ High all cancer, bladder, breast, colorectal, and lung cancer mortality rates ○ High diabetes mortality rate ○ High alcohol-related and liver disease mortality rates ○ High mortality rates among females for suicides and motor vehicle accidents ○ High smoking related neoplasm mortality among males | <ul style="list-style-type: none"> • Low hospital admission rates for ambulatory care sensitive conditions among youth and adults 18-44. • Low hospitalization rates for cardiovascular disease including CHF, AMI and Stroke • Low hospitalization rate for bronchitis/asthma among youth (0-17) • Low hospitalization rates for mental health diagnoses. • Low mortality rate for AMI • Low mortality rates for melanoma and prostate cancers • Low infant mortality rate |
|---|--|

Note: The term high connotes a result at least 10% greater than Maine result. The term low connotes a result at least 10% less than the Maine result. Highest and 2nd highest are based on comparisons between Maine counties.

Additional detail on indicators and data sources can be found in full report – Appendix 9: Detailed Data Sources

| | Oxford | Maine | Maine Counties Source |
|--|----------|-----------|-----------------------------------|
| DEMOGRAPHICS | | | |
| Total Population | 56,608 | 1,319,691 | 2008 Census Estimates |
| Median Annual Household Income (to 2008) | \$38,987 | \$46,807 | ME SPO Data Center |
| % of Labor Force Unemployed | 11% | 7.8% | ME Dept Labor |
| % Population Not Attaining H.S. Diploma (>25 yr) | 18% | 15% | 2000 Census |
| % Population on Medicaid (all ages) | 30% | 23% | 2004 CMS, HRSA Area Resource File |
| % Population Under the Age of 18 | 21% | 21% | 2008 Census Estimates |
| % Population Age 65 and Over | 16% | 15% | 2008 Census Estimates |
| % Uninsured Non-Elderly Adults (Ages 18-65) | 18% | 16% | Household Survey |
| HEALTH STATUS | | | |
| % Health Fair to Poor | 23% | 15% | Household Survey |
| % 11+ Days Lost due to Poor Mental or Physical Health | 11% | 8% | Household Survey |
| % 3+ Chronic Conditions | 17% | 13% | Household Survey |
| Wellness Categories: | | | |
| %Well | 32% | 34% | Household Survey |
| %At Risk for Future Medical Problems | 7% | 8% | Household Survey |
| %Some Health Problems | 32% | 36% | Household Survey |
| %Not Well | 29% | 23% | Household Survey |
| ACCESS TO CARE | | | |
| % Without Usual Source of Primary Care (Males) | 16% | 18% | Household Survey |
| % Without Usual Source of Primary Care (Females) | 12% | 8.3% | Household Survey |
| % Named hospital or ER as usual source of care | 2.7% | 1.9% | Household Survey |
| % Not Having a Checkup Within the Past 2 yrs (Males) | 17% | 15% | Household Survey |
| % Not Having a Checkup Within the Past 2 yrs (Females) | 6.6% | 6.4% | Household Survey |
| % Received Flu Shot or Mist past 12 months | 40% | 42% | Household Survey |
| % Ever Received Pneumococcal Vaccine (Age 65+) | 80% | 73% | Household Survey |
| % Needed Medical Care But Could not Afford it: Past Year | 8.9% | 6.5% | Household Survey |
| % No Dental Visit in Past 2 Years | 34% | 24% | Household Survey |
| ED Visits per 100,000 population | 52,871 | 47,665 | MHDO Hosp ED |
| Ages 65+ | 55,633 | 49,497 | MHDO Hosp ED |
| Hospitalizations per 100,000 Population | 12,484 | 12,076 | MHDO Hosp Inpatient |
| Ages 65+ | 31,433 | 31,396 | MHDO Hosp Inpatient |
| QUALITY/EFFECTIVENESS | | | |
| Ambulatory Care Sensitive Condition (ACSC), Hospital Admission Rate (Overall PQI*) | 915 | 967 | MHDO Hosp Inpatient |
| Ages 0-17 | 107 | 191 | MHDO Hosp Inpatient |
| Ages 18-44 | 190 | 234 | MHDO Hosp Inpatient |
| Ages 45-64 | 652 | 707 | MHDO Hosp Inpatient |
| Ages 65+ | 3,922 | 4,166 | MHDO Hosp Inpatient |
| Ambulatory Care Sensitive Condition (ACSC), ED Visit Rate (Overall PQI*) | 3,227 | 3,073 | MHDO Hosp Inpatient |
| Ages 0-17 | 2,043 | 1,994 | MHDO Hosp Inpatient |
| Ages 18-44 | 2,890 | 2,868 | MHDO Hosp Inpatient |
| Ages 45-64 | 2,411 | 2,374 | MHDO Hosp Inpatient |
| Ages 65+ | 7,101 | 6,375 | MHDO Hosp Inpatient |

All rates are per 100,000 population (based on US Census estimates 2008) unless otherwise noted

| | | Oxford | Maine | Maine Counties Source |
|------------------------------|--|--------|-------|-----------------------|
| CARDIOVASCULAR HEALTH | | | | |
| Risk Factors | % Current Smokers (Age 18+) | 25% | 22% | Household Survey |
| | % Sedentary Lifestyle (measured by no physical activity) | 24% | 21% | Household Survey |
| | % Overweight (Ages 18+) | 36% | 37% | Household Survey |
| | % Obesity (Ages 18+) | 33% | 28% | Household Survey |
| Disease Prevalence | % High Cholesterol | 35% | 29% | Household Survey |
| | % High Blood Pressure | 31% | 30% | Household Survey |
| | % Heart Disease | 6.9% | 6.3% | Household Survey |
| Management | Congestive Heart Failure, Hospital Admissions | 255 | 283 | MHDO Hosp Inpatient |
| | AMI, Hospital Admission Rate | 135 | 211 | MHDO Hosp Inpatient |
| | Ages 45-64 | 97 | 157 | MHDO Hosp Inpatient |
| | Ages 65+ | 631 | 1,037 | MHDO Hosp Inpatient |
| | Cerebrovascular Disease (stroke), Hospital Admission Rate | 141 | 149 | MHDO Hosp Inpatient |
| | CABG, Hospital Admission Rate | 68 | 62 | MHDO Hosp Inpatient |
| | % Having Cholesterol Checked within the past year (Ages 21+) | 67% | 63% | Household Survey |
| | % Smokers advised to quit smoking in the past yr. | 72% | 72% | Household Survey |
| Quality/Effectiveness | AMI, Mortality Rate | 36 | 45 | ODRVS Mortality |
| | Ages 65+ | 188 | 232 | ODRVS Mortality |
| | Cerebrovascular Disease (stroke), Mortality Rate | 55 | 49 | ODRVS Mortality |
| | Ages 65+ | 310 | 294 | ODRVS Mortality |
| | Heart Disease, Mortality Rate | 203 | 202 | ODRVS Mortality |
| | Ages 65+ | 1,052 | 1,101 | ODRVS Mortality |

All rates are per 100,000 population (based on US Census estimates 2008) unless otherwise noted

Overall PQI = methodology based on AHRQ Prevention Quality Indicators using 13 identified conditions

| | | Oxford | Maine | Maine Counties Source |
|---------------------------|--|----------------|--------------|-----------------------|
| RESPIRATORY HEALTH | | | | |
| | % Current Smokers (Male) | 29% | 23% | Household Survey |
| | % Current Smokers (Female) | 20% | 20% | Household Survey |
| | % Former Smokers | 30% | 31% | Household Survey |
| Disease Prevalence | % Current Asthma (Ages 18+) | 11% | 10% | Household Survey |
| | % Ever Asthma (Ages 0-17) | 6.9% | 6.1% | Household Survey |
| | % COPD | 6.7% | 4.2% | Household Survey |
| | Lung and Broncus Cancer, Males, Incidence Rate | 123 | 105 | ME CDC Cancer Reg |
| | Lung and Broncus Cancer, Females, Incidence Rate | 91 | 86 | ME CDC Cancer Reg |
| Management | % Received Flu Shot or Mist past 12 months | 40% | 42% | Household Survey |
| | % Ever Received Pneumoccal Vaccine (Ages 65+) | 80% | 73% | Household Survey |
| | Bronchitis and Asthma, Hospital Admission Rate Ages 65+ | 87 100 | 87 114 | MHDO Hosp Inpatient |
| | Bronchitis and Asthma, ED Visit Rate Ages 65+ | 987 559 | 988 632 | MHDO Hosp ED |
| | COPD, Hospital Admission Rate | 401 | 284 | MHDO Hosp Inpatient |
| | COPD, ED Visit Rate Ages 65+ | 1,306 2,531 | 998 1,914 | MHDO Hosp Inpatient |
| | Pneumonia, Hospital Admission Rate Ages 65+ | 465 2,105 | 326 1,402 | MHDO Hosp Inpatient |
| | Pneumonia, ED Visit Rate Ages 65+ | 531 1,235 | 505 1,053 | MHDO Hosp Inpatient |
| | Emphysema, Hospital Admission Rate Ages 65+ | 26 100 | 23 79 | MHDO Hosp Inpatient |
| | % Current Smokers advised to quit smoking in the past year | 72% | 72% | Household Survey |
| | % Current smokers tried to quit in past year | 61% | 54% | Household Survey |
| | % Current smokers ever used Maine Tobacco Quitline | 22% | 19% | Household Survey |
| | Lung Cancer, Mortality Rate (Males) | 88 | 78 | ODRVS Mortality 07-09 |
| | Lung Cancer, Mortality Rate (Females) | 73 | 61 | ODRVS Mortality 07-09 |
| | COPD, Mortality Rate (Ages 65+) | 406 | 332 | ODRVS Mortality 07-09 |
| | Pneumonia, Mortality Rate (Ages 65+) | 89 | 103 | ODRVS Mortality 07-09 |
| | Smoking-Related Neoplasms, Mortality Rate (Males) | 254 | 205 | ODRVS Mortality 07-09 |
| | Smoking-Related Neoplasms, Mortality Rate (Females) | 163 | 150 | ODRVS Mortality 07-09 |

All rates are per 100,000 population (based on US Census estimates 2008) unless otherwise noted

| | | Oxford | Maine | Maine Counties Source |
|---------------------------|--|--------|-------------------|-----------------------|
| CANCER HEALTH | | | | |
| | All Cancers, Incidence Rate | 684 | 629 | ME CDC Cancer Reg |
| | Bladder, Incident Rate | 40 | 35 | ME CDC Cancer Reg |
| | Female Breast Cancer, Incidence Rate | 178 | 162 | ME CDC Cancer Reg |
| | Female Cervix Uteri, Incidence Rate | 3.5 | 7.3 | ME CDC Cancer Reg |
| | Colorectal, Incidence Rate | 69 | 62 | ME CDC Cancer Reg |
| | Lung and Bronchus Cancer, Incidence Rate | 107 | 95 | ME CDC Cancer Reg |
| | Melanoma, Incidence Rate | 24 | 26 | ME CDC Cancer Reg |
| | Male Prostate, Incidence Rate | 184 | 187 | ME CDC Cancer Reg |
| | | | | |
| Management / Patient Care | % Reported Mammogram past year (40+) | 70% | 69% | Household Survey |
| | % Stage Female Breast, Local | 67% | 66% | ME CDC Cancer Reg |
| | % Stage Female Breast, Distant | 3.2% | 3.8% | ME CDC Cancer Reg |
| | | | | |
| | % Reported Pap Smear past 2 years | 68% | 70% | Household Survey |
| | % Stage Cervix Uteri Female, Local | 100% | 52% | ME CDC Cancer Reg |
| | % Stage Cervix Uteri Female, Distant | 0% | 14% | ME CDC Cancer Reg |
| | | | | |
| | % Reported Blood Stool Test Past Year (Age 50+) | 27% | 20% | BRFSS 2006/2008 |
| | % Reported Having Sigmoid/Colonoscopy Past 5 Yrs (Age 50+) | 61% | 63% | Household Survey |
| | % Stage Colorectal, Local | 28% | 47% | ME CDC Cancer Reg |
| | % Stage Colorectal, Distant | 36% | 17% | ME CDC Cancer Reg |
| | | | | |
| | % Stage Lung and Brunchus Male, Local | 16% | 16% | ME CDC Cancer Reg |
| | % Stage Lung and Brunchus Male, Distant | 51% | 50% | ME CDC Cancer Reg |
| | % Stage Lung and Brunchus Female, Local | 18% | 21% | ME CDC Cancer Reg |
| | % Stage Lung and Brunchus Female, Distant | 58% | 47% | ME CDC Cancer Reg |
| | | | | |
| | % Reported Prostate Exam (PSA test) past 2 yrs (males Age 50+) | 72% | 69% | Household Survey |
| | % Reported Digital Rectal Exam past 2 years (males Age 50+) | 66% | 68% | Household Survey |
| % Stage Prostate, Local | 84% | 76% | ME CDC Cancer Reg | |
| % Stage Prostate, Distant | 1.3% | 3.8% | ME CDC Cancer Reg | |
| | | | | |
| Quality/Effectiveness | All Cancers, Mortality Rate | 277 | 234 | ODRVS Mortality |
| | Bladder, Mortality Rate | 9.4 | 7.5 | ODRVS Mortality |
| | Female Breast Cancer, Mortality Rate | 38 | 28 | ODRVS Mortality |
| | Female Cervix Uteri, Mortality Rate | 2.3 | 2.5 | ODRVS Mortality |
| | Colorectal, Mortality Rate | 24 | 21 | ODRVS Mortality |
| | Lung, Mortality Rate | 80 | 69 | ODRVS Mortality |
| | Melanoma, Mortality Rate | 1.8 | 3.6 | ODRVS Mortality |
| | Male Prostate, Mortality Rate | 17 | 23 | ODRVS Mortality |
| | | | | |

All rates are per 100,000 population (based on US Census estimates 2008) unless otherwise noted

| | | Oxford | Maine | Maine Counties Source |
|---|--|--------|-------|-----------------------|
| DIABETES HEALTH | | | | |
| Disease Prevalence | % Diagnosed Diabetes | 12% | 10% | Household Survey |
| | Ages 18-44 | 5.4% | 2.9% | Household Survey |
| | Ages 45-64 | 12% | 13% | Household Survey |
| | Ages 65+ | 26% | 21% | Household Survey |
| Management | % Reported hemoglobin A1c measurement (at least once) in past year (Age 18+) | 78% | 89% | Household Survey |
| | % Reported pupil dilation eye exam in past yr (age 18+) | 84% | 76% | Household Survey |
| | % Reported foot examination in past yr (Age 18+) | 67% | 78% | Household Survey |
| | % Reported ever taken diabetes self management course (Age 18+) | 56% | 54% | Household Survey |
| | Diabetes, Hospital Admission Rate | 84 | 79 | MHDO Hosp Inpatient |
| | Ages 18-44 | 92 | 75 | MHDO Hosp Inpatient |
| | Ages 45-64 | 52 | 75 | MHDO Hosp Inpatient |
| | Ages 65+ | 172 | 150 | MHDO Hosp Inpatient |
| | Diabetes Short-term Complications, ACSC ED Visit Rate | 17 | 9 | MHDO Hosp Inpatient |
| | Diabetes Long-term Complications, ACSC ED Visit Rate | 125 | 111 | MHDO Hosp Inpatient |
| Diabetes Uncontrolled, ACSC ED Visit Rate | 11 | 11 | | |
| Quality | Diabetes, Mortality Rate | 34 | 26 | ODRVS Mortality |
| | Ages 65+ | 162 | 126 | ODRVS Mortality |

All rates are per 100,000 population (based on US Census estimates 2008) unless otherwise noted

| | | Oxford | Maine | Maine Counties Source |
|---------------------------|--|--------|--------------|-----------------------|
| MENTAL HEALTH | | | | |
| Risk Factors | % 11+ Days Mental Health Not Good | 14% | 11% | BRFSS 2008 + 2009 |
| | Ages 65+ | 5.2% | 6.1% | BRFSS 2008 + 2009 |
| | % needed, but did not get, mental health treatment in past 12 months | 6.0% | 4.8% | Household Survey |
| Disease Prevalence | % receiving outpatient mental health treatment in past 12 mos | 13% | 11% | Household Survey |
| | % At Risk for Clinical Depression Based on MHI5 (18+) | 11.9% | 7.2% | Household Survey |
| | % Diagnosed Depression (ever, 18+) | 22% | 22% | Household Survey |
| | % Current Depression (18+) | 16% | 15% | Household Survey |
| | % Diagnosed Other Psychiatric Disorder (ever, 18+) | 12% | 13% | Household Survey |
| | % Developmental Delay/Learning Disability (Ages 0-17) | 4.7% | 4.5% | Household Survey |
| Management | Psychoses Hospital Admission Rate | 572 | 578 | MHDO Hosp Inpatient |
| | Ages 65+ | 255 | 246 | MHDO Hosp Inpatient |
| | Senility and Organic Mental Disorders, Hospital Admission Rate | 5.3 | 8.6 | MHDO Hosp Inpatient |
| | Ages 65+ | 33 | 50 | MHDO Hosp Inpatient |
| | Major Depressive Disorder, Hospital Admission Rate | 145 | 157 | MHDO Hosp Inpatient |
| | Ages 0-17 | 70 | 85 | MHDO Hosp Inpatient |
| | Ages 18-64 | 188 | 196 | MHDO Hosp Inpatient |
| | Ages 65+ | 78 | 93 | MHDO Hosp Inpatient |
| | Bipolar Disorder, Hospital Admission Rate | 306 | 280 | MHDO Hosp Inpatient |
| | Ages 65+ | 89 | 88 | MHDO Hosp Inpatient |
| | Schizophrenia, Hospital Admission Rate | 84 | 114 | MHDO Hosp Inpatient |
| | Ages 65+ | 11 | 39 | MHDO Hosp Inpatient |
| | Anxiety, Hospital Admission Rate | 189 | 269 | MHDO Hosp Inpatient |
| | Ages 65+ | 54 | 70 | MHDO Hosp Inpatient |
| | Senility and Organic Mental Disorders, ED Rate | 37 | 28 | MHDO Hosp ED |
| | Major Depressive Disorder, ED Rate | 72 | 109 | MHDO Hosp ED |
| | Bipolar Disorder, ED Rate | 145 | 166 | MHDO Hosp ED |
| | Schizophrenia, ED Rate | 75 | 70 | MHDO Hosp ED |
| Anxiety Disorder, ED Rate | 1,763 | 1,618 | MHDO Hosp ED | |
| | Suicide, Mortality Rate (Males) | 20 | 23 | ODRVS Mortality |
| | Suicide, Mortality Rate (Females) | 6.9 | 5.4 | ODRVS Mortality |

All rates are per 100,000 population (based on US Census estimates 2008) unless otherwise noted

| | | Oxford | Maine | Maine Counties Source |
|---|---|--------|------------------|-----------------------|
| SUBSTANCE ABUSE | | | | |
| Prevalence | % Chronic Heavy Drinking - Past Month | 7.8% | 6.4% | BRFSS 2008 & 2009 |
| | Ages 65+ | 5.1% | 4.5% | BRFSS 2008 & 2009 |
| | % Binge Drinking -Past Month | 16% | 15% | BRFSS 2008 & 2009 |
| | Ages 18-44 | 25% | 26% | BRFSS 2008 & 2009 |
| | % Ever diagnosed with Substance Abuse Problem | 4.1% | 4.7% | Household Survey |
| | % Current Substance Abuse Problem | 0.9% | 1.5% | Household Survey |
| | % Overdose Past 12 mos (Households) | 1.5% | 0.9% | Household Survey |
| | % have used any street drugs in past 30 days | 7.2% | 5.2% | Household Survey |
| % have used any prescription drugs for non-prescribed purpose in past 30 days | 0.9% | 1.8% | Household Survey | |
| Management | Substance Abuse, Hospital Admission Rate | 359 | 379 | MHDO Hosp Inpatient |
| | Ages 65+ | 116 | 149 | MHDO Hosp Inpatient |
| | Acute Alcohol-Related Mental Disorders, Hospital Admission Rate | 22 | 50 | MHDO Hosp Inpatient |
| | Ages 65+ | 22 | 19 | MHDO Hosp Inpatient |
| | Alcohol-Related Psychoses, Hospital Admission Rate | 168 | 174 | MHDO Hosp Inpatient |
| | Ages 65+ | 28 | 52 | MHDO Hosp Inpatient |
| | Acute Drug-Related Mental Disorders, Hospital Admission Rate | 18 | 39 | MHDO Hosp Inpatient |
| | Ages 65+ | 0 | 6.5 | MHDO Hosp Inpatient |
| | Drug-Related Psychoses, Hospital Admission Rate | 152 | 117 | MHDO Hosp Inpatient |
| | Ages 65+ | 66 | 72 | MHDO Hosp Inpatient |
| | Acute Alcohol-Related Mental Disorders, ED Rate | 135 | 131 | MHDO Hosp ED |
| | Alcohol-Related Psychoses, ED Rate | 26 | 28 | MHDO Hosp ED |
| | Acute Drug-Related Mental Disorders, ED Rate | 250 | 297 | MHDO Hosp ED |
| | Drug-Related Psychoses, ED Rate | 40 | 57 | MHDO Hosp ED |
| | Alcohol-Related Mortality Rate (Males) | 25 | 19 | ODRVS Mortality |
| | Alcohol-Related Mortality Rate (Females) | 16 | 10 | ODRVS Mortality |
| | Alcohol Liver Disease, Mortality Rate | 18 | 11 | ODRVS Mortality |
| | Motor Vehicle Accidents, Mortality Rate (Males) | 23 | 21 | ODRVS Mortality |
| | Motor Vehicle Accidents, Mortality Rate (Females) | 12 | 8.0 | ODRVS Mortality |

All rates are per 100,000 population (based on US Census estimates 2008) unless otherwise noted

| | | Oxford | Maine | Maine Counties Source |
|--------------------------------|--|--------|--------------|-----------------------|
| REPRODUCTIVE HEALTH | | | | |
| | 2 or more sex partners in past yr (ages 18-34) | 19% | 16% | Household Survey |
| | % used condom last time had sex (ages 18-34) | 39% | 35% | Household Survey |
| | Teen Birth Rate (10-17yrs) Per 1,000 Females | 3.3 | 4.1 | ODRVS Birth |
| Management | High Risk Pregancy, Hospital Admission Rate (10-44 year old females) | 477 | 360 | MHDO Hosp Inpatient |
| | C-Section Rate per 100 births | 30 | 30 | ODRVS Birth |
| | % Adequate Prenatal Care (of live births) | 92% | 91% | ODRVS Birth |
| | % Inadequate Prenatal Care (of live births) | 2.5% | 2.5% | ODRVS Birth |
| | % Low Birthweight (<2500 grams) | 6.1% | 6.4% | ODRVS Birth |
| | % Prematurity (< 37 weeks) | 8.2% | 8.7% | ODRVS Birth |
| | Infant Mortality Rate (deaths to infants from birth through 364 days of age) per 1,000 live births | 4.2 | 5.5 | ODRVS Mortality |
| | Neonatal Mortality Rate (deaths to infants under 28 days) per 1,000 live births | 3.0 | 3.9 | ODRVS Mortality |
| | CHILD/YOUTH HEALTH | | | |
| | % Seriously Considered Suicide | 15% | 14% | MIYHS 2009 |
| | % Current Smoker (Past Month) (Grade 9-12) | 19% | 20% | MIYHS 2009 |
| | % Current Smokeless Tobacco User | 11% | 9.5% | MIYHS 2009 |
| | % Alcohol Use (Past Month) (Grade 9-12) | 34% | 35% | MIYHS 2009 |
| | % Binge Drink (5+ in a row) Past Month) (Grade 9-12) | 22% | 21% | MIYHS 2009 |
| | % Marijuana Use (Past Month) (Grade 9-12) | 25% | 24% | MIYHS 2009 |
| | % Sniffed Glue or Other Inhalant (Past Month) (Grade 9-12) | 10% | 9% | MIYHS 2009 |
| | % Regular Physical Activity (at least 60 min on 5 of last 7 dys) | 34% | 39% | MIYHS 2009 |
| | % Consume fruits and vegetables 5 or more times/day | 14% | 15% | MIYHS 2009 |
| | Teen Birth Rate (10-17yrs) Per 1,000 Female Population | 3.3 | 4.1 | ODRVS Birth |
| Prevalance | % Ever Been Diagnosed with Asthma (0-17) parental report | 6.9% | 6.1% | Household Survey |
| | % Overweight/Obesity Problem (0-17) parental report | 3.7% | 2.0% | Household Survey |
| | % Overweight (Grade 9-12) | 17% | 14% | MIYHS 2009 |
| | % Obese (Grade 9-12) | 16% | 13% | MIYHS 2009 |
| | % with developmental delay or learning disability (0-17) parental report | 4.7% | 4.5% | Household Survey |
| Management | ACSC, ED Rate - Overall PQI (Ages 0-17) | 2,043 | 1,994 | MHDO Hosp Inpatient |
| | ACSC, Hospital Admission Rate - Overall PQI (Ages 0-17) | 107 | 191 | MHDO Hosp Inpatient |
| | Asthma and Bronchitis, Hospital Admission Rate (Ages 0-17) | 161 | 196 | MHDO Hosp Inpatient |
| | Pneumonia, Hospital Admission Rate (Ages 0-17) | 128 | 132 | MHDO Hosp Inpatient |
| | Psychoses Hospital Admission Rate (Ages 0-17) | 503 | 484 | MHDO Hosp Inpatient |
| | Major Depressive Disorder, Hospital Admission Rate (Ages 0-17) | 70 | 85 | MHDO Hosp Inpatient |
| | Bipolar Disorder, Hospital Admission Rate (Ages 0-17) | 395 | 353 | MHDO Hosp Inpatient |
| | Asthma and Bronchitis, ED Rate (Ages 0-17) | 1,285 | 1,145 | MHDO Hosp ED |
| Pneumonia, ED Rate (Ages 0-17) | 564 | 516 | MHDO Hosp ED | |

All rates are per 100,000 population (based on US Census estimates 2008) unless otherwise noted

| | | Oxford | Maine | Maine Counties Source |
|----------------------------------|---|--------|-------|----------------------------|
| ORTHOPEDICS | | | | |
| | % Diagnosed Arthritis | 35% | 32% | BRFSS 2007 & 2009 |
| | Ages 65+ | 62% | 59% | BRFSS 2007 & 2009 |
| | Hip Procedures, Hospital Admission Rate | 120 | 93 | MHDO Hosp Inpatient |
| | Ages 65+ | 493 | 452 | MHDO Hosp Inpatient |
| | Head Brain Injury, Hospital Admission Rate | 72 | 57 | MHDO Hosp Inpatient |
| | Ages 65+ | 210 | 180 | MHDO Hosp Inpatient |
| INFECTIOUS DISEASE | | | | |
| | HIV/AIDS, Hospital Admissions Rate | 7.1 | 5.1 | MHDO Hosp Inpatient |
| | Chronic Hepatitis C, Number of Case Reports* | 35 | 1453 | ME CDC Infect Disease 2007 |
| | Sexually Transmitted Disease Incidence Rate: | | | |
| | Gonorrhea | 11 | 7.3 | ME CDC Infect Disease 2008 |
| | Chlamydia | 155 | 197 | ME CDC Infect Disease 2008 |
| INTIMATE PARTNER VIOLENCE | | | | |
| | % Ever physically hurt by Intimate Partner | 15% | 12% | Household Survey |
| | % Past yr physical violence or unwanted sex from Intimate Partner | 1.2% | 1.2% | Household Survey |

All rates are per 100,000 population (based on US Census estimates 2008) unless otherwise noted

* A Hepatitis C case report is defined as the presence of any positive serologic marker for Hepatitis C infection. State cases include reports where no county data was available so Maine total exceeds sum of counties.